

The effect of highly structured cosmic magnetic fields on ultra-high energy cosmic ray propagation

Gustavo A. Medina-Tanco^{1,2}

1. Instituto Astronômico e Geofísico, University of São Paulo, Brasil

gustavo@iagusp.usp.br

2. Dept. of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK

Received _____; accepted _____

ABSTRACT

The possibility that the magnetic field is strongly correlated with the large-scale structure of the universe has been recently considered in the literature. In this scenario the intergalactic magnetic field has a strong (μG) regular component spanning tens of Mpc but localized in sheets and filaments, while the vast voids in between are almost free of magnetic field. If true, this could have important consequences on the propagation of ultra-high energy cosmic rays, and severely affect our capacity of doing astronomy with charged particles. A quantitative discussion of these effects is given in the present work.

Subject headings: Cosmic Rays — large-scale structure — magnetic fields

1. Introduction

Recently, Ryu, Kang and Biermann (1998) proposed a large-scale scenario in which cosmic magnetic fields are clumped inside cosmological sheets and filaments in much the same way as the luminous matter does. Furthermore, their simulations suggest that the IGMF presents also a remarkable degree of alignment along cosmological structures for several Mpc. For such a model to be able to account for the existing rotational measure of extragalactic sources (e.g., Kronberg 1994), the intensity of the IGMF must be relatively high inside cosmological structures ($\sim 1\mu\text{G}$) and essentially negligible in the vast voids surrounding those structures.

This is very different from the common view in which the IGMF is roughly uniform in intensity, scaling smoothly with luminous matter density (e.g., $B_{IGM} \propto n^{0.2-0.3}$ - Vallée 1997, Medina Tanco 1997b), while the reversal scale is varied accordingly to account for the observed rotational measure. In the latter scenario, the resulting space acquires a cell-like structure (e.g., Medina Tanco, Gouveia Dal Pino and Horvath 1997b). The IGMF is more evenly distributed and its intensity ranges from $\text{few} \times 10^{-10}$ G to $\text{few} \times 10^{-9}$ G. UHECR particles random walk from their sources to the detector through this IGMF configuration producing relative small intrinsic angular error boxes (e.g., Medina Tanco, Gouveia Dal Pino and Horvath 1997b; Medina Tanco 1997b,c; Medina Tanco 1998). With high enough statistics, there is then hope for a future UHECR-astronomy.

In fact, large earthbound projects, like HiRes and Auger, and the OWL twin satellites are being proposed having such UHECR astronomy among their main priorities. However, an IGMF structure as proposed by Ryu and collaborators can, if correct, have profound consequences in our understanding of the problem of UHECR propagation in the intergalactic medium and on the kind of science one can expect to achieve with the new generations of detectors.

This work addresses the problem in a simplified, yet illustrative way. The UHECR emission coming from a flat cosmological structure, representative of a number of walls that can be encountered in the local universe (see, for example, Fairall 1998 and references there in), is analyzed. The dependence of flux, angular deflection, traversed path and other relevant parameters is studied as a function of the relative orientation between the observer, the cosmological structure where the sources are embedded and the IGMF. The effects on the observed chemical composition are also briefly discussed.

2. Numerical calculations and discussion of results

Ryu and co-workers model for the IGMF is remarkable from the UHECR point of view for at least three characteristics: (a) the high intensity of the IGMF inside walls and filaments, (b) its laminar structure and large-scale order and (c) the marked intensity contrast between the IGMF inside and outside walls separated by a relatively thin interface.

In order to analyze the effect of a large scale ordering of the intergalactic magnetic field (IGMF) on the observed sky in UHECR, an isolated wall immersed in a surrounding void is simulated. In this simplified scheme the wall is represented as a disk-like slab of 5 Mpc thickness and 20 Mpc radius. The slab is located in the x-z plane and centered in the coordinate's origin (see figure 1).

The space is permeated by a magnetic field with regular component oriented along the z-axis. The magnitude of the regular component inside the wall is constant and equal to B_{wall} , decaying to a very low value, B_{void} , inside the void. The transition between B_{wall} and B_{void} is exponential and takes place over a distance Δy measured perpendicularly to the wall. In order to break the unrealistic homogeneity of the field, a random component, B_{RND} , superimposed on the regular component, B_{reg} , is assumed. B_{RND} is characterized

by an amplitude $\eta = (B_{RND}/B_{reg})$, power law spectrum $\propto \kappa^{-\xi}$, with $\xi = 5/3$, minimum wavenumber $k_{min} = 2\pi/5 \text{ Mpc}^{-1}$ and maximum wavenumber $k_{max} = 2\pi/100 \text{ kpc}^{-1}$.

UHECR particles are injected isotropically with power law spectrum in total energy, $dN/dE \propto E^{-2}$, at 100 sources randomly distributed inside the volume of the wall. The particle sources have all the same integrated luminosity. No sources are considered in the surrounding void.

In what follows only protons are taken into account and energy losses due to redshift, pair production and photo-pion production in interactions with the cosmic microwave background radiation field are included as given by Berezhinsky and Grigor'eva (1988).

In figures 2 to 4 the results for $B_{wall} = 0.1\mu\text{G}$, $B_{void} = 10^{-10} \text{ G}$, $\eta = 0.3$ and $\Delta y = 5 \text{ Mpc}$ are shown. Note that, the value chosen for the magnetic field inside the wall is an order of magnitude below the values actually estimated by Ryu, Kang and Biermann (1998). After normalizing their simulations by the present limit in rotation measure (Kronberg 1994), they obtain $\sim 1\mu\text{G}$ on large-scales outside galaxy clusters.

For illustrative purpose, figure 2 shows the projections of the orbits of some individual particles onto the x-y plane. The general direction of the magnetic field (the regular component) is oriented perpendicular to the plane of the figure. All the particles were injected with $E = 10^{20} \text{ eV}$ at the same source in the origin of coordinates, forming an angle of 5° with the x-y plane. The only thing that varies from run to run is the azimuthal angle at injection, measured on the x-y plane with respect to, say the y-axis. It can be seen that the dynamics of the UHECR is completely different from a random walk and that a large-scale pattern of flow is to be expected. First of all, we see that particles tend to drift along the surface of the wall as soon as they reach the interface where the magnetic field decreases to its lower void value. This combines with a very efficient transport of the particle's guiding centers along the z-axis, to give a pattern of UHECR flow that is mostly

confined to the wall and preferentially directed along the general direction of the magnetic field. Furthermore, those particles that escape from the surface of the wall tend to do so in a particular direction that depends on the side of the wall that is being considered. Particles escaping towards $y > 0$ tend to fly in the $x > 0$ direction, while the opposite occurs in the $y < 0$ side of the wall. The particular injection point chosen for this example is near enough to the border of the wall to allow the escape of most of the cosmic rays produced there out to the surrounding void. Nevertheless, this escape is not isotropic. Furthermore, particles originated in sources located further inside the wall, will be mainly constrained to travel along the IGMF. This implies that a strong anisotropy must be expected in the UHECR emission originated from inside the wall. Consequently, the effects of a large-scale regular component of the IGMF in the observational properties of UHECR should be very dependent on the relative orientation between the ordered magnetic field inside the structure and the position of the detector.

The latter is clearly illustrated in figure 3, where several average quantities are shown for UHECR impinging a spherical surface of 20 Mpc radius centered in the origin of coordinates. This spherical boundary completely encloses the wall, intersecting its borders at the equator (see the sketch in figure 1). A spherical coordinate system is chosen in which the y-axis defines the polar axis and the z-axis is the origin for the longitudes, ϕ , measured on the plane of the wall (z-x). Latitudes, θ , are measured from the plane of the wall. The sphere is projected onto the plane of the page using an Aitoff equal area projection (Greisen 1993) in which the wall is the horizontal central band. The center of the plot is the point where the z-axis intersects the sphere ($\phi = 0^\circ$, $\theta = 0^\circ$) and so, in that region, the magnetic field emerges perpendicularly to the plane of the figure. The UHECR were isotropically injected with a power law energy spectrum ($dN/dE \propto E^{-2}$ and $E > 4 \times 10^{19}$ eV) at 100 sources randomly located inside the wall and, as previously mentioned, all the relevant energy losses were taken into account during propagation.

Figure 3.a shows the logarithm of the flux of UHECR (in arbitrary units). It can be seen that the charged particle flux emerging from the wall is very anisotropic, varying in almost three orders of magnitude over 4π sr. The flow of particles in the system, however, shows a definite pattern. Most of the UHECR flow along the regular component of the IGMF, producing the maxima observed near $(\phi = 0^\circ, \theta = 0^\circ)$ and $(\phi = \pm 180^\circ, \theta = 0^\circ)$. The relatively few protons ejected into the surrounding void, fly away in opposite directions above and below the plane of the wall, producing an uneven (and antisymmetric) coverage in each hemisphere. In particular, the emergent flux is a minimum when observing along the polar axis, perpendicular to the wall (i.e., from $\theta = 90^\circ$). The drift perpendicular to the IGMF direction combines with the flow along the IGMF to produce minima at $(\phi = \pm 90^\circ, \theta = 0^\circ)$, i.e., at the border of the slab but perpendicular to the magnetic field. Therefore, the emission of UHECR from an active large-scale structure can be either considerably amplified or severely hampered depending on the relative position of the observer. Broadly speaking, a boost is obtained along the IGMF and a drop in intensity in the direction perpendicular to the IGMF; a wall can be invisible for an observer located along a perpendicular through the center of the wall.

Figure 3.b shows the deflection between the arrival direction of a typical particle and the direction to the source where it originated. It can be seen that the deflection angles are very large and so any directional information is lost, except for observers looking at the wall flat on. Nevertheless, even in the latter situation, the expected error boxes are $10 - 20^\circ$ for most orientations. This could seriously hinder any possibility of astronomy with UHECR.

Figure 3.c shows the average deflection between the injection direction of a UHECR at the source and its arrival direction at the detector. Again, the expected angles are very large, around 90° for most of the sky, indicating that any such information is lost. Therefore, if acceleration mechanisms that require beaming are involved, it would be hopeless to

look for some correlation between γ and UHECR counterparts of a given event (e.g., a Gamma Ray Burst - c.f. Stanev, Schaefer and Watson 1996). This can represent a test to the hypothesis of high intensity large-scale IGMF confined to thin sheets. If both, γ and UHECR are received in correlation from a single source known to have beaming (e.g., a blazar) inside a wall, then the IGMF in the wall is most likely highly randomized and well below the $0.1\mu\text{G}$ scale.

In figure 3.d, the path lengths in Mpc traversed by the particles from the source to the sphere are displayed. Exception made of two spots antisymmetrically located at each side of the wall, the paths of the UHECR are much larger than the actual distance to their sources. This has important consequences regarding UHECR composition: if the primaries are heavy nuclei, they will more likely arrive as protons regardless the position of the detector, even for the new lower estimates of the cosmic infrared background (Stecker 1998, Malkan and Stecker 1998; see, however, Epele and Roulet 1998).

Figures 4.a-c show the Aitoff projections of the sky in UHECR originated inside the wall (roughly a hemisphere), as reconstructed by observers located in points a, b and c of figure 1 respectively. Logarithmic fluxes are represented in the contour plots and different normalizations are used for each one of the three figures. If the observer is located at the border of the wall and the large-scale IGMF is aligned with the zenithal direction (figure 4.a), the wall is visible as a huge twisted structure covering most of the sky. The rotation of the image in UHECR with respect to the wall is an indicative of the sense of the IGMF: pointing towards the observer if counter-clockwise, pointing away from the observer in the opposite case. On the other hand, the image of the wall is completely different when the zenith of the observer (located at the border of the wall and in its central plane), runs perpendicular to the IGMF (figure 4.b). In this case there is a deep minimum right at the zenith bounded by two parallel laminar structures that run roughly along the wall. Figure

4.a or 4.b (depending on the relative orientation of the IGMF) should be representative of the Centaurus wall. This feature runs edge on along the supergalactic plane (de Vaucouleurs 1956) dominating the Southern Hemisphere sky, and will be in full view of the 10^3 km^2 Auger observatory. Figure 4.c corresponds to an observer outside the wall and watching at it flat on. The image in UHECR is in this case a pair of blobs filling most of the sky, and separated by a pronounced minimum that cuts the wall in the middle perpendicularly to the IGMF direction. Furthermore, the resolution of the sky maps 4.a-c is approximately 6° and, for that resolution, no individual sources are detected.

It should be noted that the IGMF intensity assumed in the calculations presented above is one order of magnitude below the upper limit quoted by Ryu, Kang and Bierman (1998). The present analysis is therefore a conservative one in that context. However, the implications of such IGMF topology for the analysis of both existing and proposed UHECR experiments data must be seriously considered.

3. Conclusions

The quantitative results presented heretofore are model dependent and so are only intended as illustrative. Nevertheless, it has been shown that a relatively well ordered IGMF ($B_{RND}/B_{reg} \sim 0.3$) compressed to high intensities ($\sim 0.1 \mu\text{G}$) inside cosmological walls and almost negligible inside voids, dramatically changes the picture of UHECR propagation with respect to what has been normally assumed in the literature so far. In particular:

- 1 The UHECR flux leaving a wall can vary by 2 or 3 orders of magnitude over 4π , making any conclusion very dependent on the relative orientation of the flattened structure containing the sources, the IGMF inside it and the observer.
- 2 Except for privileged observers located near the central perpendicular to the wall,

directional information is mostly erased, hampering source identification.

- 3 Any information regarding the injection direction of the UHER is lost, which means that it should be impossible to detect γ and UHECR counterparts of a same bursting event if beaming is involved in the production of the primaries.
- 4 The paths traversed by the particles in their way to the detector are so large that probably no heavy primary would survive, even for the new lower estimates of the infrared background (Malkan and Stecker 1998, Stecker 1998).
- 5 The incoming flux produces a rather flat, featureless contribution at the observer's sky. In fact, at a $\sin 5^\circ$ resolution, a density of point-like sources as low as $\sim 10^{-2}$ Mpc^{-3} , gives no signal that could be identified as an individual source on the observed sky facing the wall.
- 6 The best place to test the correctness of this kind of IGMF model is the Southern Hemisphere. There, the huge Centaurus wall (home to the Great Attractor and Abell 3627) dominates the sky running roughly coincident with the Supergalactic plane and extending out to probably 6000 km/sec (~ 90 Mpc for $h = 0.65$). An additional observational bonus of the Centaurus wall is the fact that it not only seen edge on but also bordered by the very large local void. Fortunately, this large-scale structure will be in the field of view of the 10^3 km^2 southern site of the Auger experiment.

This work was done with the partial support of the Brazilian agency FAPESP.

Figure Captions

Figure 1: Schematic representation of the model used in the simulations. The horizontal slab is the wall containing randomly located UHECR point sources. It is there that the IGMF reaches its highest intensity $\sim 0.1\mu\text{G}$. The IGMF has two components. The regular component is parallel to the z-axis everywhere, its intensity is high and constant inside the wall and decays exponentially through an interface to a very low value inside the surrounding void. The random component has a power law spectrum and scales everywhere as $\eta = B_{RND}/B_{reg} = \text{const}$. Protons are injected at the sources inside the slab with a power law ($dN/dE \propto E^{-2}$ for $E > 4 \times 10^{19}$ eV) spectrum and propagated through the system until they reach the spherical border. This surface is the site of the different observers considered in the work. Several magnitude measured over this surface will be Aitoff projected in figures 3a-d, while figures 4a-c show the hemisphere of the sky facing the wall as seen by observers located in points a, b and c respectively.

Figure 2: Trajectories of several test protons injected at 10^{20} eV at origin of coordinates projected onto the x-y plane. The regular component of the magnetic field is perpendicular to the plane of the figure. All the protons were injected at an angle of 5° with respect to the x-y plane, and differ only in the azimuthal injection angle. See the text for further details.

Figure 3: Aitoff projections of: (a) the logarithm of the UHECR flux (arbitrary units), (b) deflection angle (deg) between arrival direction and true source position, (c) angle between the arrival direction and the injection direction (deg) and (d) path traversed by the UHECR in Mpc as measured over the spherical border enclosing the system (see figure 1).

Figure 4: Aitoff projection of the hemisphere of the sky facing the wall as seen by observers located in positions labeled as a, b and c in figure 1. The quantity plotted is the logarithm of UHECR flux and the normalization used is different for each case.

REFERENCES

- Berezinsky V. S. and Grigor'eva S. I., 1988, *Astron. Astrophys.*, **199**, 1.
- de Vaucouleurs G., 1956, *Vistas in astronomy*, **2**, 1584.
- Epele L. N. and Roulet E., 1998, *astro-ph/9806251*
- Fairall A., 1998, *Large-Scale Structures in the Universe*, John Wiley & Sons, England.
- Greisen E. W., 1993, *AIPS Memo* 46.
- Kronberg P. P., 1994, *Rep. Prog. Phys.*, **57**, 325.
- Malkan M. A. and Stecker F. W., 1998, *Ap. J.*, **496**, 13.
- Medina Tanco G. A., Gouveia Dal Pino E. M. and Horvath J. E., 1997a, *ApJ.*, in press
- Medina Tanco G. A., Gouveia Dal Pino E. M. and Horvath J. E., 1997b, *Astroparticle Phys.*, **6**, 337.
- Medina Tanco G. A. 1997a, *Proc. 25th Int. Cosmic Ray Conf.*, South Africa, **4**, 485.
- Medina Tanco G. A. 1997b, *Proc. 25th Int. Cosmic Ray Conf.*, South Africa, **4**, 477.
- Medina Tanco G. A. 1997c, *Proc. 25th Int. Cosmic Ray Conf.*, South Africa, **4**, 481.
- Medina Tanco G. A., 1998, *Ap. J. Lett.*, **495**, L71-L74, 1998.
- Ryu D., Kang H. and Biermann P. L., 1998, *A.& A.*, **335**, 19.
- Stanev T., Schaefer R. and Watson A. A., 1996, *Astroparticle Phys.*, **5**, 75.
- Stecker F. W., 1998, *Phys. Rev. Lett.*, **80**, 1816.
- Vallée J. P., 1997, *Fundamentals of Cosmic Phys.*, **19**, 1.

This figure "figure1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9808073v1>

This figure "figure2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9808073v1>

This figure "figure3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9808073v1>

This figure "figure4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9808073v1>